

References

- Becker, J., Nakajima, M., & Wu, W. F. (2018). Factors contributing to medial tibial stress syndrome in runners: A prospective study. *Medicine & Science in Sports & Exercise*, 50(10), 2092–2100. <https://doi.org/10.1249/mss.0000000000001674>
- Bhusari, N., & Deshmukh, M. (2023). Shin Splint: A review article. *Cureus*. <https://doi.org/10.7759/cureus.33905>
- Bramah, C., Preece, S. J., Gill, N., & Herrington, L. (2019). A 10% increase in step rate improves running kinematics and clinical outcomes in runners with patellofemoral pain at 4 weeks and 3 months. *The American Journal of Sports Medicine*, 47(14), 3406–3413. <https://doi.org/10.1177/0363546519879693>
- C. Madden, C., Putukian, M., C. McCarty, E., & C. Young, C. (2017). Netter's Sports Medicine 2nd edition. *The Bangkok Medical Journal*, 13(02), 113–113. <https://doi.org/10.31524/bkkmedj.2017.09.020>
- Franklyn, M. (2015). Aetiology and mechanisms of injury in medial tibial stress syndrome: Current and future developments. *World Journal of Orthopedics*, 6(8), 577. <https://doi.org/10.5312/wjo.v6.i8.577>
- Lee, I., Jeon, H. G., Ha, S., Jeong, H., & Lee, S. Y. (2025). How medial tibial stress syndrome is affected by alignment, range of motion, strength, and gait biomechanics: A systematic review and meta-analysis. *Journal of Sport Rehabilitation*, 34(2), 134–155. <https://doi.org/10.1123/jsr.2024-0031>
- McClure, C. J. (2023). Medial tibial stress syndrome. *StatPearls*. <https://doi.org/10.5040/9781350969100>

Milner, C. E., Hamill, J., & Davis, I. (2007a). Are knee mechanics during early stance related to tibial stress fracture in runners? *Clinical Biomechanics*, 22(6), 697–703.

<https://doi.org/10.1016/j.clinbiomech.2007.03.003>

Milner, C. E., Hamill, J., & Davis, I. (2007b). Are knee mechanics during early stance related to tibial stress fracture in runners? *Clinical Biomechanics*, 22(6), 697–703.

<https://doi.org/10.1016/j.clinbiomech.2007.03.003>

Moen, M. H., Bongers, T., Bakker, E. W., Zimmermann, W. O., Weir, A., Tol, J. L., & Backx, F. J. (2012). Risk factors and prognostic indicators for medial tibial stress syndrome.

Scandinavian Journal of Medicine & Science in Sports, 22(1), 34–39.

<https://doi.org/10.1111/j.1600-0838.2010.01144.x>

Naderi, A., Fallah Mohammadi, M., Heidaralizadeh, A., & Moen, M. H. (2025). Effects of integrating lower-leg exercises into a multimodal therapeutic approach on medial tibial stress syndrome management among recreational runners: A randomized controlled study. *Orthopaedic Journal of Sports Medicine*, 13(2).

<https://doi.org/10.1177/23259671241311849>

Napier, C., MacLean, C. L., Maurer, J., Taunton, J. E., & Hunt, M. A. (2018). Kinetic risk factors of running-related injuries in female recreational runners. *Scandinavian Journal of Medicine & Science in Sports*, 28(10), 2164–2172.

<https://doi.org/10.1111/sms.13228>

Nigg, B., Baltich, J., Hoerzer, S., & Enders, H. (2015). Running shoes and running injuries: Mythbusting and a proposal for two new paradigms: ‘preferred movement path’ and ‘comfort filter.’ *British Journal of Sports Medicine*, 49(20), 1290–1294.

<https://doi.org/10.1136/bjsports-2015-095054>

- Ohta-Fukushima, M., Mutoh, Y., Takasugi, S., Iwata, H., & Ishii, S. (2002). Characteristics of stress fractures in young athletes under 20 years. *J Sports Med Phys Fitness*.
- Saad, M. A., Jamal, J. M., Aldhafiri, A. T., & Alkandari, S. A. (2025). Medial tibial stress syndrome: A scoping review of epidemiology, biomechanics, and risk factors. *Cureus*. <https://doi.org/10.7759/cureus.81463>
- Tenforde, A. S., Sayres, L. C., McCurdy, M. L., Collado, H., Sainani, K. L., & Fredericson, M. (2011). Overuse injuries in high school runners: Lifetime prevalence and prevention strategies. *PM&R*, 3(2), 125–131. <https://doi.org/10.1016/j.pmrj.2010.09.009>
- Vincent, H. K., & Vincent, K. R. (2022). Healthy running habits for the Distance Runner: Clinical Utility of the American College of Sports Medicine Infographic. *Current Sports Medicine Reports*, 21(12), 463–469. <https://doi.org/10.1249/jsr.0000000000001019>
- Warden, S. J., Davis, I. S., & Fredericson, M. (2014). Management and prevention of bone stress injuries in long-distance runners. *Journal of Orthopaedic & Sports Physical Therapy*, 44(10), 749–765. <https://doi.org/10.2519/jospt.2014.5334>
- Warden, S. J., Edwards, W. B., & Willy, R. W. (2021). Optimal load for managing low-risk tibial and metatarsal bone stress injuries in runners: The science behind the clinical reasoning. *Journal of Orthopaedic & Sports Physical Therapy*, 51(7), 322–330. <https://doi.org/10.2519/jospt.2021.9982>
- Warden, S. J., Hoenig, T., Sventeckis, A. M., Ackerman, K. E., & Tenforde, A. S. (2022). Not all bone overuse injuries are stress fractures: It is time for updated terminology. *British Journal of Sports Medicine*, 57(2), 76–77. <https://doi.org/10.1136/bjsports-2022-106112>
- Wille, C. M., Lenhart, R. L., Wang, S., Thelen, D. G., & Heiderscheit, B. C. (2014). Ability of sagittal kinematic variables to estimate ground reaction forces and joint kinetics in

running. *Journal of Orthopaedic & Sports Physical Therapy*, 44(10), 825–830.

<https://doi.org/10.2519/jospt.2014.5367>

Willy, R. W., Buchenic, L., Rogacki, K., Ackerman, J., Schmidt, A., & Willson, J. D. (2015). In-field gait retraining and mobile monitoring to address running biomechanics associated with tibial stress fracture. *Scandinavian Journal of Medicine & Science in Sports*, 26(2), 197–205. <https://doi.org/10.1111/sms.12413>

Yates, B., & White, S. (2004). The incidence and risk factors in the development of medial tibial stress syndrome among naval recruits. *The American Journal of Sports Medicine*, 32(3), 772–780. <https://doi.org/10.1177/0095399703258776>